

Gardiner's phonon for Bose-Einstein Condensation: A physical realization of the q -deformed Boson

Chang-Pu Sun^{ab}, Sixia Yu

Institute of theoretical Physics, Academia Sinica, Beijing 100080, P.R. China

Yi-Bao Gao

Department of Physics, Beijing Polytechnic University, Beijing 100022, P.R. China

The Gardiner's phonon presented for a particle-number conserving approximation method to describe the dynamics of Bose-Einstein Condensation (BEC) (C.W. Gardiner, Phys. Rev. A 56, 1414 (1997)) is shown to be a physical realization of the q -deformed boson, which was abstractly developed in quantum group theory. On this observation, the coherent output of BEC atoms driven by a radio frequency (r.f) field is analyzed in the viewpoint of a q -deformed Fock space. It is illustrated that the q -deformation of bosonic commutation relation corresponds to the non-ideal BEC with the finite particle number N of condensated atoms. Up to order $1/N$, the coherent output state of the untrapped atoms *minimizes* the uncertainty relation like a coherent state does in the ideal case of BEC that N approaches infinity or $q = 1 - 2/N$ approaches one.

I. INTRODUCTION

Since the introduction of the concept of q -deformed boson [1] by different authors independently in the development of the quantum group theory [2-4], there have been many efforts of finding its physical realizations [5,6], e.g., fittings the deformed spectra of rotation and oscillation for molecules and nuclei [7,8]. In our opinion, those investigations can be regarded as merely phenomenological because a q -deformed structure is postulated in advance without giving it a microscopic mechanism. In this paper it will be shown that a physical and natural realization of the q -deformed boson is provided by the phonon operators, which was presented recently by Gardiner [10] for the description of Bose-Einstein condensation (BEC).

As well known, to deal with the dynamics of Bose-Einstein condensated gas the Bogoliubov approximation in quantum many-body theory is an efficient approach, in which the creation and annihilation operators for condensated atoms are substituted by c -numbers. One shortcoming of this method is that the total atomic particle-number may not be conserved after the approximation. Or a symmetry may be broken. To remedy this default, Gardiner suggested a modified Bogoliubov approximation by introducing phonon operators which conserve the total atomic particle number N and obey the bosonic commutation relations in case of $N \rightarrow \infty$. In this sense this phonon operator approach gives an elegant infinite atomic particle number approximation theory for BEC taking into account the conservation of the total atomic number.

What will be investigated here is the case that the total atomic particle number N is very large but not infinite. That is, we shall consider the effects of order $o(1/N)$. And we shall focus on an algebraic method of treating the effects of finite condensated particle number in atomic BEC. As it turns out, the commutation rela-

tions for the Gardiner's phonon operators will no longer obey the commutation relation of the Heisenberg-Weyl algebra but the q -deformed bosonic commutation relation

$$[b, b^\dagger]_q \equiv bb^\dagger - qb^\dagger b = 1, \quad (1)$$

where the deformation constant q depends on the total atomic particle number. By using this q -deformed boson algebra we will discuss the effects of the finiteness of the particle number on the output of MIT experiment of BEC.

This paper is organized as follows. In sec.II after the reformulation of the Gardiner's modified Bogoliubov method for the two-level atoms, the q -deformed commutation relation for the Gardiner's phonon operator is derived in case of large but finite particle number. The excitation of the Gardiner's phonons is also considered for small N . In sec.III, the Hamiltonian describing the coherent output in MIT experiment for atomic laser [13] is discussed in terms of the q -deformed algebra. When $q = 1$ or $N \rightarrow \infty$ it describes the coherent output of BEC from a vacuum to a coherent states. For general q , or finite but large condensated atomic particle number, it results in a nonlinear dynamic Heisenberg equation for the q -deformed bosonic operators. Its solution at the first order of $1/N$ is analyzed. In sec.IV dressed phonon, a generalization of the Gardiner's phonon in case of quantized radiation field, is discussed and it satisfies also a q -deformed bosonic algebra for a large photon number and atomic particle number. Finally there are some concluding remarks in the last section.

II. Q -DEFORMED BOSONIC ALGEBRA FOR THE GARDINER'S PHONON

Gardiner's starting point to introduce the phonon operators is to consider a system of the weakly interacting

Bose gas. Without losing generality, we consider for the moment a classical radiation field interacting with two-level atoms with $b_e^\dagger$ and b_e denoting the creation and annihilation operators for the atoms in the untrapped state and $b_g^\dagger$ and b_g for the creation and annihilation operators of the atoms in the trapped ground state. The simplified Hamiltonian for the MIT experiment reads

$$H = \hbar\omega_e b_e^\dagger b_e + \hbar g(t) b_e^\dagger b_g + \text{h.c.}, \quad (2)$$

where $g(t)$ is a time-dependent coupling coefficient for the classical sweeping r.f. field coupled to those two states with level difference $\hbar\omega_e$. Note that the total atomic particle number $N = b_e^\dagger b_e + b_g^\dagger b_g$ is conserved. For convenience we define $\eta = 1/N$ for large particle number.

In the thermodynamical limit $N \rightarrow \infty$, the Bogoliubov approximation is usually applied, in which the ladder operators $b_g^\dagger, b_g$ of the ground state are replaced by a c -number $\sqrt{N_c}$, where N_c is the average number of the inital condensated atoms. As a result Hamiltonian Eq.(2) becomes a forced harmonic oscillator with the external force term $\hbar\sqrt{N_c}g(t)b_e^\dagger + \text{h.c.}$ Obviously, it will drive the atom in the untrapped excited state from a vacuum state to a coherent state and thus lead to a coherent output of BEC atoms in the propagating mode. In this case the condensated atoms are initially described by a coherent state [13].

However, this approximation destroys a symmetry of the Hamiltonian Eq.(2), i.e., the conservation of the total particle number is violated because of $[N, H_b] \neq 0$. According to Gardiner, the phonon operators are defined as:

$$b = \frac{1}{\sqrt{N}} b_g^\dagger b_e, \quad b^\dagger = \frac{1}{\sqrt{N}} b_g b_e^\dagger. \quad (3)$$

These operators act invariantly on the subspace V^N spanned by bases $|N; n\rangle \equiv |N - n, n\rangle$ ($n = 0, 1, \dots, N$), where Fock states

$$|m, n\rangle = \frac{1}{\sqrt{m!n!}} b_e^{\dagger m} b_g^{\dagger n} |0\rangle$$

with $m, n = 0, 1, 2, \dots$ span the Fock space H_{2b} of a two mode boson.

A straightforward calculation leads to the following commutation relation between the phonon operator and its Hermitian conjugate:

$$[b, b^\dagger] = 1 - \frac{2}{N} b_e^\dagger b_e = f(b^\dagger b; \eta), \quad (4)$$

with $f(x; \eta) = \sqrt{1 + 2(1 - 2x)\eta + \eta^2} - \eta$. Keeping only the lowest order of η for a very large total atomic particle number, the commutator above becomes

$$[b, b^\dagger] = 1 - 2\eta b^\dagger b \quad (5a)$$

or

$$[b, b^\dagger]_q = 1, \quad (5b)$$

with $q = 1 - 2\eta$. This is exactly a typical q -deformed commutation relation. As $N \rightarrow \infty$ or $q \rightarrow 1$, the usual commutation relation of Heisenberg-Weyl algebra is regained.

Since the Gardiner's phonon excitation for BEC can be mathematically understood in terms of q -deformed boson system, the algebraic method previously developed in quantum group theory can be of use for the construction of a q -deformed Fock space. The existence of a vacuum state $|0\rangle$ such that $b|0\rangle = 0$ is guaranteed by the fundamental structure of the q -deformed boson algebra [14]. Starting with the vacuum, from relation

$$bb^{\dagger n} = q^n b^{\dagger n} b + \langle n \rangle b^{\dagger n-1}, \quad (6)$$

where the q -deformed c -number $\langle n \rangle = n - n(n-1)\eta$ is up to the first order of η and approaches to n when $N \rightarrow \infty$, the q -deformed Fock space for the Gardiner's phonon excitation of BEC is spanned by bases

$$|n\rangle = \frac{1}{\sqrt{\langle n \rangle!}} b^{\dagger n} |0\rangle \quad (7)$$

with $\langle n \rangle! = \langle n \rangle \langle n-1 \rangle \cdots \langle 1 \rangle$ ($n = 0, 1, \dots, N$). The action of operators b and $b^\dagger$ are then represented on the deformed Fock space as:

$$b^\dagger |n\rangle = \sqrt{\langle n+1 \rangle} |n+1\rangle, \quad b |n\rangle = \sqrt{\langle n \rangle} |n-1\rangle \quad (8)$$

the actions of the creation and annihilation operators b and $b^\dagger$ coincide with their direct actions on the subspace V^N of the Fock space of a two-mode boson on which the total particle number is conserved.

Actually, the q -deformed vacuum state is physically a two-mode boson state $|N; 0\rangle$ and it is trivially annihilated by the phonon operator b . So the vacuum state can be regarded as a case that all bosons condensed in the ground state and tends to form a BEC. By mapping back to two-mode boson space, the q -deformed Fock state $|n\rangle \sim |N; n\rangle$ implies a phonon excitation that n bosons are created in the propagating mode by destroying n bosons in trapped mode. This kind of fundamental process of phonon excitation essentially guaranteed the particle number conservation.

In the above discussion about the Gardiner's phonon excitation, we have linearized commutator $h \equiv f(b^\dagger b; \eta)$ so that a q -deformed commutation rule was obtained. Essentially this linearization establishes a physical realization of the q -deformed algebra. However, if the total particle number N is not large enough, then h can not be approximated by a linear function. From the commutation relations between h and $b, b^\dagger$

$$[h, b^\dagger] = -\frac{2}{N} b^\dagger, \quad [h, b] = \frac{2}{N} b, \quad (9)$$

we see that the algebra of Gardiner's operators is a rescaling of algebra $su(2)$ with factor N . When $N \rightarrow \infty$, it approaches the Heisenberg-Weyl algebra because h approaches identity. Its representation space is a finite dimensional space is found to be spanned by bases given in

Eq.(7) with $\langle n \rangle = n - n(n-1)\eta$ being exact. It is interesting to notice that $\langle n \rangle$ coincide with the q -deformed number in the first order of η . The action of those creation and annihilation operators have the same form. On this finite dimensional space we have $b|0\rangle = 0$ and $h|0\rangle = |0\rangle$ together with $b^\dagger|N\rangle = 0$. It is worthy to point out that the last equality can not be composed to the usual Heisenberg-Weyl algebra because such a constrain will make it is impossible for the creation operator to be the Hermitian conjugate to the annihilation operator. Physically the usual boson algebra and its q -deformation with q being not a root of unity describe systems without particle number conservation such as the black body radiation, but this finite dimensional representation provide us with a suitable algebraic framework to treat the dynamic process where the particle number is conserved.

III. THE COHERENT OUTPUT OF BEC AS A Q -DEFORMED BOSONIC EXCITATION

In this section we shall apply the concept of the q -deformed boson algebra for the Gardiner's phonon excitation to the theoretical analysis of the MIT experiment for the coherent output of the BEC atoms from a trap.

By denoting $N_e = b_e^\dagger b_e$, in terms of the deformed boson operators the Hamiltonian can be rewritten as:

$$H_q = \hbar\omega_e N_e + \hbar\sqrt{N} (g(t)b^\dagger + \text{h.c.}). \quad (10)$$

For an ideal BEC with infinite atoms condensated in the trapped state, the q -deformed bosonic commutation relation becomes the usual commutation relation $[b, b^\dagger] = 1$ of HW algebra. Together with commutator $[N_e, b] = -b$, a linear Heisenberg equation

$$\dot{b} = -i\omega_e b - i\mu(t) \quad (11)$$

is acquired where $\mu(t) = g(t)\sqrt{N}$ is finite as N tends to infinity. Its solution can be easily found to be

$$b(t) = b(0)e^{-i\omega_e t} + \beta(t) \quad (12)$$

with a non-vanishing expectation value in the vacuum state:

$$\beta(t) = \langle 0|b(t)|0\rangle = -i \int_0^t e^{-i\omega_e(t-t')} \mu(t') dt'. \quad (13)$$

Therefore when the BEC is initially at a Fock state, one obtains a coherent state output of the untrapped atoms. This is just a direct manifestation of the quantum coherence through a factorization $\langle b^\dagger(t)b(t) \rangle = \langle b^\dagger(t) \rangle \langle b(t) \rangle$. In fact for infinite N , the average phonon number $\langle b^\dagger(t)b(t) \rangle$ approach the average atomic number $b_e^\dagger b_e$ of outputting in the untrapped mode.

Further, to consider the quantum coherence of the outputting atoms represented by phonon operators $b(t)$, we

evaluate the variances of the Hermitian quadrature phase operators [16]

$$X_1(t) = \frac{b(t) + b^\dagger(t)}{\sqrt{2}}, \quad X_2(t) = \frac{b(t) - b^\dagger(t)}{i\sqrt{2}}. \quad (14)$$

One can easily find that

$$\Delta^2 X_{1,2}(t) = \langle 0|X_{1,2}^2|0\rangle - \langle 0|\Delta X_{1,2}|0\rangle^2 = \frac{1}{2} \quad (15)$$

are time-independent constants that minimize the uncertainty product $\Delta^2 X_1(t)\Delta^2 X_2(t) = 1/4$. It reflects that the outputted atoms are just in a coherent state. This judgements for the case with infinite N can be similarly used to analyses the influence of q -deformation with finite atomic number on the dynamics of outputting BEC.

In practice condensated atomic number N cannot be infinite and therefore the Bogoliubov approximation can not work well and so an approximation method with higher accuracy that for the generic Bogoliubov approximation or its Gardiners modification. And the q -deformed boson algebra disscussed in the previous section provides us with a algebraic structure of treating this correction of finite atomic particle number.

Up to the approximation of order η , as discussed previously, the dynamics of Gardiner's phonon excitation for coherent output of BEC is determined by the q -deformed boson commutation relation. The corresponding Heisenberg equation

$$\dot{b}(t) = -i\omega_e b(t) - i\mu(t) + 2i\eta\mu(t)b^\dagger(t)b(t) \quad (16)$$

is then a nonlinear one. It is should be remarked here that that the effective coupling $\mu(t) = g(t)\sqrt{N}$ is finite in the thermodynamical limit of an infinite number of atoms in an infinite volume but with the density fixed, i.e., $n = \lim_{N,V \rightarrow \infty} (N/V)$ is finite. This is because the coupling constant $g(t)$ between the r.f. pulse and atoms must be proportional to the inverse of the square root of the the effective mode volume V . So we have $\mu(t) \propto \sqrt{N/V}$, the square root of the density of the condensated atoms in the thermodynamical limit. For this reason the nonlinear terms in Eq.(16) can be handled as a perturbation in comparison with other terms. Otherwise, if $\mu(t)$ were not fixed as N approaches infinity, the non-liner term should not be regarded as a perturbation due to its square amplification of perturbation. The thermodynamical limit is our key point to deal with the dynamics of q -deformation for the coherent output of the BEC atoms with a perturbation method.

Becasue of the nature of the deformed bonon, in general, it is only neccissary to solve this nonlinear Heisenberg equation Eq.(16) according to the first order of η by perturbation. In fact if the annihilation operator at time t is expanded as

$$b(t) = b_0 e^{-i\omega_e t} + \beta(t) + \eta b_1(t) \quad (17)$$

with $b_0 = b(0)$, one obtains immediately

$$b_1(t) = \xi(t)b_0 + \beta^2(t)e^{i\omega_e t}b_0^\dagger - 2\beta(t)b_0^\dagger b_0 + \alpha(t), \quad (18a)$$

$$\alpha(t) = 2i \int_0^t e^{-i\omega_e(t-t')} \mu(t') |\beta(t')|^2 dt', \quad (18b)$$

$$\xi(t) = 2i \int_0^t e^{-i\omega_e t} \mu(t') \beta^*(t') dt'. \quad (18c)$$

As a result the average outputted atomic number $\langle N_e(t) \rangle$ in the untrapped excited state with finite condensated BEC particle number correction is given by

$$\langle N_e(t) \rangle = |\beta(t)|^2 + \eta \left(|\beta(t)|^4 + 2\text{Re}(\beta(t)\alpha(t)^*) \right). \quad (19)$$

The quantum fluctuation of the quadrature phase operators $X_1(t)$ and $X_2(t)$ in the final state can also be explicitly determined as:

$$\Delta^2 X_1 = \frac{1}{2} - \eta \left(|\beta(t)|^2 + \beta^2(t) + \beta^{*2}(t) \right), \quad (20a)$$

$$\Delta^2 X_2 = \frac{1}{2} - \eta \left(|\beta(t)|^2 - \beta^2(t) - \beta^{*2}(t) \right). \quad (20b)$$

Keeping the first order of η , one obtains immediately the following relation

$$\Delta^2 X_1 \Delta^2 X_2 = \frac{1}{4} - \eta |\beta(t)|^2. \quad (21)$$

The term of η reflects the deviation from the thermodynamical limit as the effect of the q -deformation with finite condensated particle number.

Since the commutation relation of quadrature phase operators $X_1(t)$ and $X_2(t)$ is deformed into

$$[X_1(t), X_2(t)] = i(1 - 2\eta b^\dagger(t)b(t)), \quad (22)$$

the corresponding Heisenberg's uncertainty relationship of those quadratures in the output taking account of finite atomic number should be:

$$\Delta^2 X_1 \Delta^2 X_2 \geq \frac{1}{4} - \eta \langle 0 | b^\dagger(t)b(t) | 0 \rangle. \quad (23)$$

Considering $\langle 0 | b^\dagger(t)b(t) | 0 \rangle = |\beta(t)|^2 + o(\eta)$ the coherent output state of the untrapped atoms *minimizes* the uncertainty relation. Because, as we known, the squeezed state and coherent state can be defined to be states minimize the uncertainty relation between the position and momentum, the output state of the atomic lase can be likened to a kind of q -deformed coherent state or even a squeezed state because fluctuations Eq.(26a) and Eq.(26b) are inequal.

IV. DRESSED PHONON OPERATOR

When the r.f field is a quantized one which is described by a single-mode boson whose annihilation operator is denoted as a , we consider the following Hamiltonian:

$$H_d = \omega_e N_e + \omega_g N_g + \omega_0 N_0 + g(a^\dagger b_g^\dagger b_e + ab_g b_e^\dagger), \quad (24)$$

where $N_0 = a^\dagger a$ is the photon number operator and g is a real coupling constant. Obviously there is a conservative quantity $\Delta = N_0 + N_e$ in addition to N . We now define the dressed phonon operators as

$$B = \frac{1}{\sqrt{N\Delta}} a^\dagger b_g^\dagger b_e, \quad B^\dagger = \frac{1}{\sqrt{N\Delta}} ab_g b_e^\dagger. \quad (25)$$

For convenience we denote $\eta_0 = 1/\Delta$ for large Δ . The dressed phonon operators satisfy the following commutation relation:

$$[B, B^\dagger] = 1 - 2(\eta_0 + \eta)N_e + \eta_0\eta(3N_e^2 - N_e). \quad (26)$$

In terms of those dressed phonon operators the Hamiltonian Eq.(24) can be rewritten as

$$H_d = \omega_g N + \omega_0 \Delta + \omega_\delta N_e + \mu_d (B + B^\dagger). \quad (27)$$

where $\mu_d = g\sqrt{N\Delta}$ and $\omega_\delta = \omega_e - \omega_g - \omega_0$.

As N (or Δ) tends to infinity the dressed phonon operators $B, B^\dagger$ and N_e will give a representation of algebra $su(2)$. As both N and Δ tends infinity one obtains Heisenberg-Weyl algebra $[B, B^\dagger] = 1$. When we keep the first order in $o(\eta)$ and $o(\eta_0)$, and considering $B^\dagger B = N_e + o(\eta + \eta_0)$, we obtain a q -deformed boson algebra

$$[B, B^\dagger]_{q_d} = 1 \quad (28)$$

with the deforming constant given by $q_d = 1 - 2(\eta + \eta_0)$.

Within this kind of approximation, the Heisenberg equation of motion satisfied by the dressed phonon is given by:

$$\dot{B} = -i\omega_\delta B - i\mu_d + 2i(\eta_0 + \eta)\mu_d B^\dagger B. \quad (29)$$

Here the coupling constant g is proportional to the inverse of the square root of volume V which makes μ_d finite as N and Δ approaches infinity. Because the evolution equation Eq.(28) is the same with Heisenberg equation Eq.(16), all the results discussed in the previous section should also be valid in case of quantized r.f. field.

V. REMARKS

To conclude this paper, we treat the same problem in the Schrödinger picture. Without the effect of q -deformation caused by finite N , the dynamic process can be described as a factorized evolution of the wave function [16]. Suppose the initial state is $|\phi(0)\rangle = |\alpha\rangle = \sqrt{N}\rangle_g \otimes |0\rangle_e$ i.e., all atoms occupied the trapped ground state with a coherent state while there is no atom in the propagating mode. Driven by a r.f. field $g(t) = g \exp(i\omega_f t)$ ($g \propto \sqrt{1/V}$) with a fixed frequency ω_f , the system will have exactly a factorized evolution wave function of the form $|\phi(t)\rangle = |\alpha\rangle_g \otimes |\beta(t)\rangle_e$ even for a finite N . Such a factorization structure of wave function first

pointed out by Mewes et al is very crucial to realize their experiment, and its dynamical origins mainly depend on the linearity of the original coupling system of b_e and b_g . The coherent state of the second component $|\beta(t)\rangle_e$ of $|\phi(t)\rangle$ means a coherent output of atoms in propagating mode in the MIT experiment. The influence of finite N on Schroedinger wave function is manifested by the outputting amplitude $|\beta(t)| \propto \sqrt{N/V} |\sin(\Omega t)|$ where $\Omega = \sqrt{(\omega - \omega_f)^2/4 + g^2}$. Because the factor $\sqrt{N/V}$ is finite and Ω tends to $|\omega - \omega_f|$ in the thermodynamics limit, the results from the Bogoliubov approximation mentioned in Sec.II is just recovered by this exact solution.

There is another nonlinear effect different from that resulting from the q -deformation with finite N . In presence of interatomic interaction in the trapped or untrapped states, which roughly is of the form $b_i^\dagger b_j^\dagger b_k b_l$, the corresponding Heisenberg equations are no longer linear and thus the factorization structure will be broken down. It is noticed that such a non-linearity is essentially different from that resulting from the q -deformation with finite N in this paper. The later originates from the subsystem isolation of a large system.

ACKNOWLEDGMENT

This work is partly supported by the NFS of China.

Townsend and W. Ketterle, Phys. Rev. Lett. 78, 583 (1997)

[14] C.P. Sun, IL. Nuuvo. Cimento B 34, 499 (1993)

[15] C.P. Sun, Y.B. Gao, H.F. Dong and S.R. Zhao Phys. Rev. E 57 (1998) 3900

[16] C.P. Sun, H. Zhan, Y.X. Miao, and J.M. Li Commun. Theoretical Phys. 161, 29 (1998)

^a Electronic address: suncp@itp.ac.cn

^b Internet www site: <http://www.itp.ac.cn/~suncp>

[1] M. Arik and D. Coon, J. Math. Phys. 17, 524 (1976)

[2] L.C. Biedenharn, J. Phys. A 22, L873 (1989)

[3] C.P. Sun and H.C.Fu, J. Phys. A 22, L983 (1989)

[4] A.J. Macfarlan, J. Phys. A 22, 4851 (1989)

[5] For a review article, see C.P. Sun and M.L. Ge *The q deformed boson realization of quantum algebra and its application to Yang-Baxter equation*, in *Quantum Group and Quantum Integrable Sstems* (World Scientific, ed. M.L. Ge) 133 (1992)

[6] O.W. Greenberg, Phys. Rev. D 43, 4111 (1991)

[7] D. Bonatsos, E.N. Argyres, S.B. Prenska, P.P. Reychev, R.R. Roussev and Y.F. Smirnov, Phys. Lett B 251, 477 (1990); J. Phys. G. 16, 137 (1990)

[8] Z. Chang, W. Chen, H.Y. Guo and H. Yang, J. Phys. A 23 4185 (1991); J. Phys. A 24 1427 (1991)

[9] J.M. Li and C.P. Sun Phys. Lett. A 180, 50 (1993)

[10] C.W. Gardiner, Phys. Rev. A 56, 1417 (1997)

[11] N.N. Bogoliubov, J. Phys. (USSR) 11, 23 (1947)

[12] A.L. Fetter and J.D. Walecka, *Quantum Theory of Many-Particle System* (McGraw-Hill, 1971)

[13] M.Mewes, M.R. Andrews, D.S. Durfee, D.M. Kurn, C.G.